



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

On the Transitive Substitution Groups whose Order is a Power of a Prime Number.

BY G. A. MILLER.

In a transitive group G of degree n , the subgroup G_1 , which contains all the substitutions of G that do not involve a given letter, is of degree $n - \alpha$ ($\alpha \geq 1$), and G_1 is one of n/α conjugate substitutions of G .^{*} Each of these subgroups is, therefore, transformed into itself by αg_1 substitutions of G , g_1 being the order of G_1 . These substitutions constitute a group G_2 of order αg_1 . When $\alpha > 1$, G_2 contains a constituent of degree α . Since each of the substitutions of G_2 that is not contained in G_1 contains the α letters which G_1 omits, and since the order of G_2 is αg_1 , it follows that the said constituent of G_2 is a regular group of order α . In what follows we shall assume that the order G is a power of a prime p^m . As the order of a subgroup G must be a power of the same prime $\alpha g_1 = p^k$; hence, $\alpha = p^k$. This result may be stated as follows:

THEOREM I.—*If the order of a transitive group is a power of a prime p^m , the subgroup formed by all its substitutions which omit a given letter omits p^k ($k \geq 1$) letters of the group.*

Any set of conjugate subgroups or substitutions of G is transformed by all the substitutions of G according to a transitive substitution group of order p^k . Hence, Theorem I includes the theorem that all the substitutions of G which transform one of these subgroups or substitutions into itself, must also transform p^k ($k \geq 1$) of its conjugates into themselves. In other words, the substitutions of G which transform one of a set of conjugate subgroups or substitutions into itself constitute a group in which p^k of these conjugates are invariant.[†] This includes the theorems. Every non-invariant subgroup or substitution of a group of order

^{*}Cf. Netto, "Theory of Substitutions," 1892, p. 84. Also Cauchy, *Comptes Rendus*, vol. 21, 1845, p. 669.

[†]Burnside, "Theory of Groups," 1897, p. 65.

p^m is transformed into itself by p^k ($k \geq 1$) of its conjugates. A group of order p^{m-1} that is contained in a group of order p^m is invariant. A group of order p^m cannot be generated by one set of its conjugate subgroups.

Let K represent the group formed by all the substitutions in the holomorph* of G which transform the substitutions of G according to its group of cogredient isomorphisms. The order of K is some power of p , and its subgroup formed by all the substitutions that omit a given letter is the group of cogredient isomorphisms of G . From the facts that each letter of this subgroup corresponds to a substitution of G ,† and that this subgroup omits p^k ($k \geq 1$) letters of G , it follows that G contains p^k invariant substitutions. Hence, the given theorem includes the important theorem, due to Sylow, that every group of order p^m contains invariant operators besides identity.

When G_1 is transitive, it must be holomorphic to the regular constituent of G_2 mentioned above, since G_2 contains at least one other subgroup which is conjugate with G_1 under G . As all of these conjugates are transitive, there can be only two of them. This is only possible when $p = 2$ and $k = \frac{m-1}{2}$. Hence, the

THEOREM II.—*If the subgroup formed by all the substitutions which omit one letter of a transitive group of order p^m is transitive, the order of the group is 2^{2n+1} , n being any integer.*

When the condition of this theorem is satisfied, G_2 is clearly the direct product of G_1 and its other conjugate. Hence, it follows from a known theorem‡ that the number of transitive substitution groups of order 2^{2n+1} , whose largest subgroups of degree lower than the degree of the group are transitive, is equal to the number of regular groups of degree 2^n .

Since each of these groups may be constructed by writing a regular group of order 2^n in two distinct sets of letters and adding to their direct product a substitution of order two which permutes the corresponding letters of its systems of intransitivity, the number of invariant operators of such a group must be the same as the number of such operators in the mentioned regular group of order 2^n . The largest Abelian subgroup that is contained in such a group is clearly the

* Bulletin of the American Mathematical Society, vol. VI, 1900, p. 396.

† Ibid., vol. V, 1899, p. 245.

‡ Quarterly Journal of Mathematics, vol. 28, 1896, p. 207. American Journal of Mathematics, vol. XXI, 1899, p. 306.

direct product of the Abelian subgroups of these regular groups of order 2^n . Hence, the transitive groups of order p^m in which the subgroup formed by all the substitutions which omit a given letter is transitive, constitute an infinite system of groups of order 2^{2^n+1} which are completely determined by the groups of order 2^n .

Suppose that G_1 contains k systems of intransitivity. The number of systems of intransitivity of G_2 is then $\leq k+1$. We shall first show that $p \leq k+1$. The largest subgroup of G which transforms G_2 into itself must transform G_1 into p^λ ($\lambda \geq 1$) of its conjugates, and hence it must contain $k+1-h$ ($p-1$)($h \geq 1$) systems of intransitivity. This proves that $p \leq k+1$. When G_1 is transitive, $k=1$ and $p=2$ as was observed above.

When $p = k+1$, the largest subgroup of G that transforms G_2 into itself is transitive. Since this transitive subgroup is of the same degree as G and contains the same subgroup that omits one letter, it must be G itself. In this case G_2 contains p similar regular constituents of order α . These results may be stated as follows:

THEOREM III.—*If the subgroup formed by all the substitutions which omit one letter of a transitive group of order p^m contains k systems of intransitivity, then $p \leq k+1$. When $p = k+1$, the transitive constituents of this subgroup are similar and regular.*

Cauchy proved that the symmetric group of degree n contains subgroups of order p^m , where $m = \varepsilon\left(\frac{n}{p}\right) + \varepsilon\left(\frac{n}{p^2}\right) + \varepsilon\left(\frac{n}{p^3}\right) + \dots$; $\varepsilon\left(\frac{a}{b}\right)$ being the largest integer which does not exceed $\frac{a}{b}$.* We proceed to determine when such a group (G) is transitive and to study some of the properties of these groups. Since the symmetric group of degree n contains all the possible substitutions in n letters, G is of degree n whenever $n \equiv 0 \pmod{p}$. The degree of each of the transitive constituents of G must be a power of p , as it is a divisor of p^m . If G were intransitive when $n = p^\beta$, we would have

$$p^{\beta_1} + p^{\beta_2} + p^{\beta_3} + \dots = p^\beta,$$

$p^{\beta_1}, p^{\beta_2}, p^{\beta_3}, \dots$ being the degrees of the transitive constituents of G . Hence, the number of constituents of lowest degree would be a multiple of p . As these constituents would all be similar, we could combine p of them and thus form a

* Cauchy, *Comptes Rendus*, vol. 21, 1845, p. 844.

transitive constituent of a larger order in the same letters. This is impossible, as $n!$ is not divisible by p^{m+1} . G must, therefore, be transitive when $n = p^\beta$.

It is clear that all the symmetric groups of degrees $p^\beta + \alpha$, $\alpha < p$ contain the same G , and that the G 's of all the symmetric groups of degrees $p^\beta + \alpha'$, $\alpha' < p^{\beta+1} - p^\beta$ are the direct products of this G and the G of the symmetric group of degree α' . That is, when α' is p^β ($p > 2$), the corresponding G is the direct product of the G of the symmetric group of degree p^β and its conjugate written in a distinct set of letters; when $\alpha' = 2p^\beta$ ($p > 3$), the corresponding G is the direct product of three such groups; when $\alpha' = lp^\beta + j$ ($p > l + 1$, $j < p^\beta$) the corresponding G is the direct product of $l + 1$ such groups and the largest group whose order is a power of p that is contained in the symmetric group of degree j . Hence, the

THEOREM IV.—*The largest group G of order p^m that is contained in the symmetric group of degree n is transitive whenever $n = p^{m'} + \alpha$, $\alpha < p$, and only then. When this condition is satisfied, G contains a subgroup of order p^{m-1} , which is the direct product of p conjugate transitive groups of order $p^{\frac{m-1}{p}}$. These transitive groups in turn contain subgroups of order $p^{\frac{m-1}{p^2}-1}$, which are the direct products of p conjugate transitive groups of order $p^{\frac{m-p-1}{p^2}}$, etc.*

COROLLARY I.—*In a group of order p^m , every subgroup whose order exceeds p^{m-n-1} ($m > p^{n-1} + p^{n-2} + \dots + 1$) is invariant or contains an invariant subgroup of order $\geq p$.*

COROLLARY II.—*The largest subgroup of order p^m that is contained in any symmetric group contains just p^γ invariant operators, γ being the number of its transitive constituents.*

This theorem was proved above. To see that it involves Corollary I it is only necessary to observe that a group of order g which contains a subgroup of order g_1 , which is not invariant nor contains an invariant subgroup of the entire group, can be represented as a transitive substitution group of degree $g \div g_1$.* Corollary II follows from the fact that each of the transitive constituents of the mentioned subgroups of order $p^{\frac{m-1}{p}}$ contains just p invariant operators.

We proceed to give a method by means of which it is possible to construct a transitive group of order p^m (m being any number greater than 2) which con-

* Dyck, *Mathematische Annalen*, vol. XXII (1883), p. 102.

tains only p invariant operators. Let H represent a regular group of order p^α which contains only p invariant operators and whose quotient group with respect to these invariant operators contains no operator whose order exceeds p , and let H_1 be the conjugate of H which is formed by all the substitutions (in the same letters as are contained in H) which are commutative with every substitution of H .* Since the quotient group of H_1 with respect to its p invariant operators contains no operator whose order exceeds p , H_1 contains a non-Abelian subgroup of order p_β which includes its p invariant operators. This subgroup and H generate a group of order $p^{\alpha+2}$ which contains only p invariant operators with respect to which its quotient group contains no operator whose order exceeds p . Since it is well known that groups of the given type exist when $\alpha = 3$ or 4 ,† it follows that they exist for every value of $\alpha > 2$.

It follows from the preceding paragraph that the only value of m for which a group of order p^m must contain more than p invariant operators is two. It is easily seen that this is also the only value of m for which every subgroup of order p^{m-2} is invariant; for if a group of order p^α contains a non-invariant subgroup of order $p^{\alpha-2}$, any direct product of which this group is a factor must have the same property. Since groups of order p^3 contain non-invariant subgroups of order p , there must be groups of order p^m (m being any integer > 2) which contain non-invariant subgroups of order p^{m-2} .

The following method may be employed to determine all the groups of order p^m , provided all the groups of order p^{m-1} are known. Suppose that a group (R) of order p^m is represented as a regular group. Any one of its subgroups (H) of order p^{m-1} contains p systems of intransitivity which are permuted according to the group of order p by the remaining substitutions of R . Hence, H may be constructed by writing after each substitution of a regular group of order p^{m-1} the same substitution in $p - 1$ distinct sets of letters.‡ All of the other substitutions of R are of the form st , where t merely interchanges the corresponding letters of the p systems of H and s transforms each one of these systems into itself.§

Since t is completely determined by H , it is only necessary to consider how s may be selected. When the substitutions of H are transformed by R

* Jordan, "Traité des substitutions," 1870, p. 60.

† Hölder, *Mathematische Annalen*, vol. XLIII, 1893, p. 410.

‡ Quarterly Journal of Mathematics, vol. XXVIII, 1896, p. 236.

§ Ibid.

according to a substitution in its group of cogredient isomorphisms, we may assume that s is commutative with each substitution of H . In this case it may evidently be assumed that s involves only the letters of the first system of intransitivity of H , for, if it were otherwise, we could transform st by a substitution which would transform H into itself and also reduce the number of letters in s . Hence, there cannot be more such groups than the number of sets of substitutions in H which are conjugate under its holomorph. This number may sometimes be reduced by the following considerations:

Let $s_1, s_2, s_3, \dots, s_p$ represent the constituents of a substitution of H , each constituent involving all the letters of one of the transitive constituents of H . From the equations

$$\begin{aligned} (s_1^{p-1}s_2s_3^2s_4^3\dots s_p^{p-2})^{-1}ts_1^{p-1}s_2s_3^2s_4^3\dots s_p^{p-2} \\ = (s_1^{p-1}s_2\dots s_p^{p-2})^{-1}ts_1^{p-1}s_2\dots s_p^{p-2}t^{-1}t = s_1^{1-p}s_2s_3s_4\dots s_pt, \end{aligned}$$

it follows that s may be so selected that it is not the p^{th} power of any substitution in the first transitive constituent of H . Since s must clearly be commutative with every substitution of H , none of the powers of a non-commutative substitution in the first transitive constituent of H is a suitable value of s . In particular, we observe that s can have only one value when H is cyclical or when it is Abelian and of type $(1, 1, 1, \dots)$. It has been assumed throughout that s' is not identity, and that R transforms H according to a substitution in its group of cogredient isomorphisms.

When some substitutions of R transform H according to a substitution which is not in its group of cogredient isomorphisms, we may write these substitutions in the form st_1t , where t_1 and t are commutative, while s is commutative with every substitution of H . As in the preceding case, we may assume that s involves only the letters of the first transitive constituent of H , and hence it is also commutative with t_1 . It is evident that t_1 may be restricted to at most one out of each conjugate set of operators of order p^s in the group of isomorphisms of H .